

TEOSYAL RHA® 1



For dynamic & delicate softening of fine lines on the face, particularly crow's feet and oral commissures.¹

¹. TEOSYAL RHA® 1 – Instructions For Use.

TEOSYAL RHA® 1

Almost
7 OUT OF 10
PATIENTS

had lasting effects
through 1 year¹

TEOSYAL RHA® 1

TECHNICAL SPECIFICITIES



Hyaluronic Acid (HA) concentration	15 mg/mL
Composition	Mix of cross-linked & non-cross-linked HA
MoD BDDE cross-linking	~1.9%
Rheological Properties	Very high Stretch
Indications¹	Superficial wrinkles: ○ Upper Face ○ Perioral: Perioral rhytids, lips and philtrum, oral commissures ○ Midface ○ Neck & décolleté
Estimated duration*	At least 12 months ²
Lidocaine	0.3%
Packaging	2x1 mL
Needle/Cannula	Needle 30G or cannula 25G/50mm

BDDE : 1,4-Butanediol Diglycidyl Ether.

*Average duration depends on several factors: patients' skin type, wrinkle severity, injection type and volume

1. TEOSYAL RHA® 1 – Instructions For Use

2. Sundaram H et al. «Global Aesthetics Consensus: Hyaluronic Acid Fillers and Botulinum Toxin Type A-Recommendations for Combined Treatment and Optimizing Outcomes in Diverse Patient Populations.» Plast Reconstr Surg, pp. 137(S), p. 1410-23, 2016.

TEOSYAL RHA® 1

RHEOLOGICAL PARAMETERS¹




- ● ● ● STRETCH
- ○ ○ ○ STRENGTH
- ● ● ● STRUCTURAL INTEGRITY

HA concentration: 15 mg/mL

RHA Collection

BY TEOXANE



TEOSYAL RHA® 1 – Instructions For Use

1. Favre J, Gallet M, Teinturier C, Teyssie P, Bourdon F. Advanced Concepts in Rheology for the Evaluation of Hyaluronic Acid-Based Soft Tissue Fillers. *Dermatol Surg.* 2021;47(5):e159-e167.

RHA® 1 = Teosyal® RHA 1. RHA® 2 = Teosyal® RHA 2. RHA® 3 = Teosyal® RHA 3. RHA® 4 = Teosyal® RHA 4



TEOSYAL RHA® 1

MDR INDICATIONS

Treatment options*



- ● ● ● STRETCH
- ○ ○ ○ STRENGTH
- ● ● ● STRUCTURAL INTEGRITY

HA concentration: 15 mg/mL



SUPERFICIAL LAYERS

○ SKIN QUALITY	✓
○ FOREHEAD LINES	✓
○ EYEBROWS	✓
○ CROWS' FEET	✓
○ PERIORAL RHYTIDS	✓
○ NECK & DÉCOLLETÉ	✓
○ NECKLINES	✓